arXiv:1211.5774v1 [math.DG] 25 Nov 2012

SOME UNSTABLE CRITICAL METRICS FOR THE $L^{\frac{n}{2}}$-NORM OF THE CURVATURE TENSOR

ATREYEE BHATTACHARYA AND SOMA MAITY

Abstract. We consider the Riemannian functional defined on the space of Riemannian metrics with unit volume on a closed smooth manifold M given by $\mathcal{R}_{\frac{n}{2}}(g) := \int_M |R(g)|^{\frac{n}{2}} dv_g$ where $R(g)$, dv_g denote the Riemannian curvature and volume form corresponding to g. We show that there are locally symmetric spaces which are unstable critical points for this functional.

1. Introduction

Let M be a closed smooth manifold of dimension $n \geq 3$ and $\mathcal{M}_1$ be the space of Riemannian metrics with unit volume on M endowed with the $C^{2,\alpha}$-topology for any $\alpha \in (0,1)$. In this paper we study the following Riemannian functional

$$\mathcal{R}_p(g) = \int_M |R(g)|^p dv_g$$

where $R(g)$ and dv_g denote the corresponding Riemannian curvature tensor and volume form, $p \in [2, \infty)$. Let $S^2(T^*M)$ be the space of symmetric two tensors on M and $\mathcal{W}$ be the subspace of $S^2(T^*M)$ orthogonal to the tangent space of the orbit of g under the action of the group of diffeomorphisms of M at g. Let H denote the Hessian of $\mathcal{R}_p$ at any critical metric. For definitions of critical metric and Hessian we refer to section 2.

Definition 1.1. Let g be a critical point for $\mathcal{R}_{p|\mathcal{M}_1}$. g is *stable* for $\mathcal{R}_p$ if there is an $\epsilon > 0$ such that for every element h in $\mathcal{W}$,

$$H(h,h) \geq \epsilon \|h\|^2 \tag{1.1}$$

where $\|.\|$ denote the L^2-norm on $S^2(T^*M)$ defined by g.

Spherical space forms are stable for $\mathcal{R}_p$ for $p \geq 2$ and hyperbolic manifolds are stable for $p \geq \frac{n}{2}$ [2]. The stability of locally symmetric spaces is not known in general. In this paper we prove the following.

Theorem 1. *Let (M,g) be an irreducible locally symmetric space of compact type. If the universal cover of M is one of the following then (M,g) is not stable for $\mathcal{R}_{\frac{n}{2}}$.*

$$SU(q)(q \geq 3), \quad Sp(q)(q \geq 2), \quad Spin(5), \quad Spin(6), \quad SU(2q+2)/Sp(q+1),$$
$$Sp(q+l)/Sp(q) \times Sp(l)(l, q \geq 1), \quad E_6/F_4, \quad F_4/Spin(9)$$

Moreover, (M,g) is a saddle point for $\mathcal{R}_{\frac{n}{2}}$.

The theorem follows by restricting H to the space of conformal variations of any irreducible symmetric space and using an estimate for the first positive eigenvalue of the Laplacian of (M,g) .

Let (M,g) be a simply connected irreducible symmetric space of compact and λ_1 and s denote its first positive eigenvalue of the Laplacian and scalar curvature of it. We prove that if $\frac{\lambda_1}{s} \geq \frac{2}{n}$

Key words and phrases. Riemannian functional, stability.

then (M,g) is stable for $\mathcal{R}_{\frac{n}{2}}$ restricted to the conformal variations of g. The above condition is also a necessary and sufficient criterion for the stability of the identity map of (M,g) as a harmonic map. In [2] the stability of the identity map of these spaces has been studied in detail. We observe that if (M,g) is not a sphere then g is stable for $\mathcal{R}_{\frac{n}{2}}$ if and only if it is stable for the identity map.

Let (M,g) be an irreducible symmetric space of compact type or a compact quotient of an irreducible locally symmetric space of non-compact type. From the proof of the theorem we observe that if (M,g) is neither one of the type in Theorem 1 then (M,g) is stable for $\mathcal{R}_p (p \geq \frac{n}{2})$ restricted to the conformal variations of (M,g).

2. PROOF

Let $\{e_i\}$ be an orthonormal basis at a point of M. $\check{R}$ is a symmetric 2-tensor defined by

$$\check{R}(x,y) = \sum R(x,e_i,e_j,e_k)R(y,e_i,e_j,e_k).$$

Let D and D^* be the Riemannian connection, its formal adjoint and s denote the scalar curvature. $d^D : S^2(T^*M) \to \Gamma(T^*M \otimes \Lambda^2 M)$ and its formal adjoint δ^D are defined by

$$\begin{aligned} d^D\alpha(x,y,z) &:= (D_y\alpha)(x,z) - (D_z\alpha)(x,y) \\ \delta^D(A)(x,y) &= \sum\{D_{e_i}A(x,y,e_i) + D_{e_i}A(y,x,e_i)\} \end{aligned}$$

where $\Lambda^2 M$ and $\Gamma(T^*M \otimes \Lambda^2 M)$ denote alternating two forms and sections of $T^*M \otimes \Lambda^2 M$. Let g_t be a one-parameter family of metrics with $\frac{d}{dt}(g_t)_{|t=0} = h$ and $T(t)$ be a tensor depending on g_t. Then $\frac{d}{dt}T(t)_{|t=0}$ is denoted by $T'_g(h)$. Define $\Pi_h(x,y) = \frac{d}{dt}D_x y_{|t=0}$ where x,y are two fixed vector fields. The suffix h will be omitted when there will not be any ambiguity. Consider any $f \in C^\infty(M)$. Note that

$$\begin{aligned} g(\Pi_{fg}(x,y),z) &= \frac{1}{2}[D_x fg(y,z) + D_y fg(x,z) - D_z fg(x,y)] \qquad (2.1) \\ &= \frac{1}{2}[df(x)g(y,z) + df(y)g(x,z) - df(z)g(x,y)] \end{aligned}$$

Let Δ denote the Laplace operator which acts on $C^\infty(M)$. We use the following definition.

$$\Delta f = -tr(Ddf)$$

$(,)$, $|.|$, $\langle,\rangle$, $\|.\|$ denote point-wise inner product, point-wise norm, global inner product and global norms induced by g.

$\nabla\mathcal{R}_p(g)$ in $S^2(T^*M)$ is called the *gradient* of $\mathcal{R}_p$ at g if for every $h \in S^2(T^*M)$,

$$\frac{d}{dt}_{|t=0}\mathcal{R}_p(g+th) = \mathcal{R}'_{p|g}.h = \langle\nabla\mathcal{R}_p(g), h\rangle$$

g is called a *critical point* for $\mathcal{R}_{p|\mathcal{M}_1}$ if the component of $\nabla\mathcal{R}_p(g)$ along the tangent space of $\mathcal{M}_1$ at g is zero. The Hessian at a critical point of $\mathcal{R}_p$ is given by

$$H(h_1,h_2) = \langle(\nabla\mathcal{R}_p)'_g(h_1), h_2\rangle \quad \forall\ h_1,h_2 \in S^2(T^*M)$$

Proposition 2.1. *Let (M,g) be a compact irreducible symmetric space and $f \in C^\infty(M)$. Then*

$$H(fg,fg) = p|R|^{p-2}[a\|\Delta f\|^2 - b\|df\|^2 + c\|f\|^2]$$

where, a, b, c is given by,

$$\begin{aligned} a &= n - 1 + (p-2)\frac{4s^2}{n^2|R|^2} \\ b &= 4(p-1)\frac{s}{n} \\ c &= (p - \frac{n}{2})|R|^2. \end{aligned}$$

Let (M, g) be a closed irreducible symmetric space and $h_1, h_2 \in S^2(T^*M)$. From [2] (4.1) we have,

$$\begin{aligned} H(h_1, h_2) \;=\; & -p|R|^{p-2}(\langle \delta^D (D^*)'_g(h_1)R, h_2\rangle + \langle D^* R'_g(h_1), d^D h_2\rangle) - p|R|^{p-2}\langle \check{R}'_g(h_1), h_2\rangle \quad (2.2) \\ & -p\langle (|R|^{p-2})'_g(h_1)R, Dd^D h_2\rangle - \frac{p}{n}|R|^2\langle (|R|^{p-2})'_g(h_1)g, h_2\rangle \\ & +\frac{1}{2}\langle (|R|^p)'_g(h_1)g, h_2\rangle + \frac{p}{n}\|R\|^p\langle h_1, h_2\rangle \end{aligned}$$

Next we compute each term of the above equation for conformal variations to obtain the Proposition.

Lemma 2.1. $\langle \delta^D (D^*)'(fg)R, fg\rangle = 4\frac{s}{n}\|df\|^2$

Proof. Let $\tilde{g}(t)$ be an one-parameter family of metrics with $\tilde{g}(0) = g$ and T be a $(0, 4)$ tensor independent of t. Expressing D^* in a local coordinate chart and differentiating it, we obtain,

$$(D^*T)(x, y, z) = -(\tilde{g}^{kj})'(D_k T)_{jxyz} + \tilde{g}^{kj}[T_{\Pi_{kj}xyz} + T_{j\Pi_{kx}yz} + T_{jx\Pi_{ky}z} + T_{jxy\Pi_{kz}}]$$

Note that, Π acting on two vector fields gives a vector field. Now evaluating $(D^*)'_g(h)(R)$ on an orthonormal basis we have,

$$(D^*)'_g(h)(R)_{jkl} = R_{\Pi_{ii}jkl} + R_{i\Pi_{ij}kl} + R_{ij\Pi_{ik}l} + R_{ijk\Pi_{il}}.$$

From the definition of d^D we have,

$$d^D fg(x, y, z) = D_y fg(x, z) - D_z fg(x, y) = df(y)g(x, z) - df(z)g(x, y)$$

Combining these two,

$$\sum (D^*)'(fg)_{jkl} d^D(fg)_{jkl} \;=\; 2\sum [R_{\Pi_{ii}jkj} + R_{i\Pi_{ij}kj} + R_{ij\Pi_{ik}j}]df_k$$

Let μ be the Einstein constant of (M, g). Now using (2.2) we have,

$$\begin{aligned} 2\sum R_{\Pi_{ii}jkj} df_k &= 2\sum g(\Pi(e_i, e_i), e_m)R(e_m, e_j, e_k, e_j)df(e_k) \qquad (2.3) \\ &= -(n-2)\mu|df|^2 \end{aligned}$$

Similarly,

$$2\sum R_{ij\Pi_{ikj}} df_k = n\mu|df|^2 \text{ and } 2\sum R_{i\Pi_{ij}kj} df_k = 2\mu|df|^2 \qquad (2.4)$$

Combining the equations (2.3) and (2.4) the proof of the lemma follows. ☐

Lemma 2.2. $\langle D^* R'_g(fg), d^D fg\rangle = -(n-1)\|\Delta f\|^2 + (n-4)\frac{s}{n}\|df\|^2$

Proof. By a simple calculation we have,

$$D^2_{x,y} fg(u, v) = Ddf(x, y)g(u, v) \qquad (2.5)$$

and

$$Dd^D fg(x, y, z, w) = Ddf(x, z)g(y, w) - Ddf(x, w)g(y, z). \qquad (2.6)$$

From [1] 1.174(c) and using (2.5) we have,

$$\begin{aligned} R'_g(fg)(x,y,z,u) &= -\frac{1}{2}[Ddf(y,z)g(x,u)+Ddf(x,u)g(y,z)-Ddf(x,z)g(y,u) \\ &\quad -Ddf(y,u)g(x,z)]+fR(x,y,z,u) \end{aligned} \tag{2.7}$$

Therefore,

$$\begin{aligned} (R'_g(fg), Dd^D fg) &= Ddf_{ik}R'_g(fg)_{ijkj} - Ddf_{il}R'_g(fg)_{ijjl} \\ &= -(n-2)|Ddf|^2 - |\Delta f|^2 - 2\mu|df|^2 \end{aligned}$$

Using Bochner-Weitzenbök formula on the space of one forms we have,

$$\Delta df = D^*Ddf + (n-1)cdf$$

Hence the lemma follows. ☐

Lemma 2.3. $\langle \check{R}'_g(fg), fg\rangle = 4\frac{s}{n}\|df\|^2 - |R^2|\|f\|^2$

Proof.

$$\check{R}_{pq} = \tilde{g}^{i_1i_2}\tilde{g}^{j_1j_2}\tilde{g}^{k_1k_2}R_{pi_1j_1k_1}R_{qi_2j_2k_2}$$

Differentiating it with respect to t and evaluating on an orthonormal basis we have,

$$\begin{aligned} (\check{R}_g.h)'_{pq} &= -h_{mn}\left(R_{pmij}R_{qnij} + R_{pimj}R_{qinj} + R_{pijm}R_{qijn}\right) \\ &\quad +(R'_g.h)_{pijk}R_{qijk} + R_{pijk}(R'_g.h)_{qijk} \end{aligned}$$

Therefore,

$$\langle \check{R}'_g(fg), fg\rangle = -3|R|^2\|f\|^2 + 2\langle (R'_g.fg), fR\rangle$$

Using (2.7) we have,

$$\begin{aligned} (\check{R}'_g(fg), R) &= \frac{1}{2}\sum[Ddf_{jk}R_{ijki} + Ddf_{il}R_{ijjl} - Ddf_{ik}R_{ijkj} - Ddf_{jl}R_{ijil}] + f|R|^2 \\ &= 2\mu\Delta f + f|R|^2 \end{aligned} \tag{2.8}$$

Hence the lemma follows. ☐

Lemma 2.4. $(|R|^p)'(fg) = 2\frac{s}{n}p|R|^{p-2}\Delta f - pf|R|^p$

Proof.

$$\begin{aligned} (|R|^p)'(fg) &= p|R|^{p-2}(R, R'_g.fg) - 2p|R|^{p-2}(\check{R}, fg) \\ &= p|R|^{p-2}(R, R'_g.fg) - 2pf|R|^{p-2}tr(\check{R}) \\ &= p|R|^{p-2}(2\mu\Delta f + f|R|^2) - 2pf|R|^p \\ &= 2\mu p|R|^{p-2}\Delta f - pf|R|^p \end{aligned}$$

☐

Using (2.6) we have,

$$(Dd^D fg, R) = 2(Ddf, r) = -2\mu\Delta f$$

Now the Proposition follows from the above lemma and equation.

Proof of Theorem 1.1: Let (M,g) is a simply connected irreducible symmetric space of compact type which is not a sphere. Then

$$R = \frac{s}{n(n-1)}I + W$$

where I is the curvature of standard sphere with sectional curvature 1 and W is the Weyl curvature of (M, g). From the above expression we have

$$\frac{s^2}{|R|^2} < \frac{2}{n(n-1)}$$

Let λ_1 be the first positive eigenvalue of the Laplacian of (M, g) and f be an eigenfunction corresponding to λ_1. Then from Proposition 2.1 we have,

$$\begin{aligned} H(fg, fg) &= s\lambda_1 p|R|^{p-2}[a(\frac{\lambda_1}{s}) - \frac{b}{s}]\|f\|^2 \\ &\leq s\lambda_1 p|R|^{p-2}[(n-1+4\frac{n-4}{n^3(n-1)})\frac{\lambda_1}{s} - 2\frac{n-2}{n}]\|f\|^2 \end{aligned}$$

From the Table A.1 and A.2 in [2] we have,

$$H(fg, fg) < 0.$$

Next, we choose a sufficiently large eigenvalue λ_i such that $a(\frac{\lambda_i}{s}) - \frac{b}{s} < 0$. Let $\tilde{f}$ be an eigenfunction corresponding to λ_i. Then we have,

$$H(\tilde{f}g, \tilde{f}g) < 0$$

This completes the proof. □

Theorem 2. *Let (M, g) be either a compact quotient of an irreducible symmetric of non-compact type or a compact symmetric space which is not one of the types in Theorem 1. (M, g) is stable for $\mathcal{R}_p$ for $p \geq \frac{n}{2}$ when it is restricted to the space of conformal variations of g.*

Proof. If (M, g) is a a compact quotient of an irreducible symmetric of non-compact type then the theorem is an immediate consequence of Proposition 2.1. Otherwise from the Table A.1 and A.2 in [2] we have $\frac{\lambda_1}{s} \geq \frac{2}{n}$. Therefore,

$$\begin{aligned} H(fg, fg) &\geq sp\lambda_1|R|^{p-2}[a(\frac{\lambda_1}{s}) - \frac{b}{s}]\|f\|^2 \\ &\geq sp\lambda_1|R|^{p-2}[(n-1)\frac{\lambda_1}{s} - 2\frac{n-2}{n}]\|f\|^2 \\ &\geq 2sp\lambda_1|R|^{p-2}[\frac{n-1}{n} - 2\frac{n-2}{n}]\|f\|^2 \\ &\geq 0 \end{aligned}$$

□

Remark : Let (M, g) be one of the critical metrics of $\mathcal{R}_p$ mentioned in Theorem 2. This is an immediate consequence of the above theorem that (M, g) is a local minimizer for $\mathcal{R}_p (p \geq \frac{n}{2})$ restricted to the space of metrics conformal to g.

REFERENCES

[1] Arthur L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3)[Results in Mathematics and Related Areas (3)], Volume **10**, Springer-Verlag, Berlin, (1987).

[2] Soma Maity, *On the stability of the L^p-norm of curvature tensor*, arXive:1201.1691[math.DG] 15 March 2012.

[2] Hajime Urakawa, *The first eigenvalue of the Laplacian for a positively curved homogeneous Riemannian manifold*, Composito Mathematica, Volume **59**, p. 57-71(1986).

Department of Mathematics, Indian Institute of Science, Bangalore-12, India
E-mail address: `somamaity@math.iisc.ernet.in, atreyee@math.iisc.ernet.in`